


RESEARCH

Open Access



AtCesA8-driven *OsSUS3* expression leads to largely enhanced biomass saccharification and lodging resistance by distinctively altering lignocellulose features in rice

Chunfen Fan^{1,2,3}, Shengqiu Feng^{1,2,3}, Jiangfeng Huang^{1,2,3}, Yanting Wang^{1,2,3}, Leiming Wu^{1,2,3}, Xukai Li^{1,2,3}, Lingqiang Wang^{1,2,3}, Yuanyuan Tu^{1,2,3}, Tao Xia^{1,2,4}, Jingyang Li^{1,3,5}, Xiwen Cai⁶ and Liangcai Peng^{1,2,3*} 

Abstract

Background: Biomass recalcitrance and plant lodging are two complex traits that tightly associate with plant cell wall structure and features. Although genetic modification of plant cell walls can potentially reduce recalcitrance for enhancing biomass saccharification, it remains a challenge to maintain a normal growth with enhanced biomass yield and lodging resistance in transgenic plants. Sucrose synthase (SUS) is a key enzyme to regulate carbon partitioning by providing UDP-glucose as substrate for cellulose and other polysaccharide biosynthesis. Although *SUS* transgenic plants have reportedly exhibited improvement on the cellulose and starch based traits, little is yet reported about *SUS* impacts on both biomass saccharification and lodging resistance. In this study, we selected the transgenic rice plants that expressed *OsSUS3* genes when driven by the *AtCesA8* promoter specific for promoting secondary cell wall cellulose synthesis in *Arabidopsis*. We examined biomass saccharification and lodging resistance in the transgenic plants and detected their cell wall structures and wall polymer features.

Results: During two-year field experiments, the selected *AtCesA8::SUS3* transgenic plants maintained a normal growth with slightly increased biomass yields. The four independent transgenic lines exhibited much higher biomass enzymatic saccharification and bioethanol production under chemical pretreatments at $P < 0.01$ levels, compared with the controls of rice cultivar and empty vector transgenic line. Notably, all transgenic lines showed a consistently enhanced lodging resistance with the increasing extension and pushing forces. Correlation analysis suggested that the reduced cellulose crystallinity was a major factor for largely enhanced biomass saccharification and lodging resistance in transgenic rice plants. In addition, the cell wall thickenings with the increased cellulose and hemicelluloses levels should also contribute to plant lodging resistance. Hence, this study has proposed a mechanistic model that shows how *OsSUS3* regulates cellulose and hemicelluloses biosyntheses resulting in reduced cellulose crystallinity and increased wall thickness, thereby leading to large improvements of both biomass saccharification and lodging resistance in transgenic rice plants.

Conclusions: This study has demonstrated that the *AtCesA8::SUS3* transgenic rice plants exhibited largely improved biomass saccharification and lodging resistance by reducing cellulose crystallinity and increasing cell wall thickness. It also suggests a powerful genetic approach for cell wall modification in bioenergy crops.

Keywords: Sucrose synthase, Transgenic rice, Biomass saccharification, Yeast fermentation, Lodging resistance, Cell wall, Cellulose crystallinity

*Correspondence: lpeng@mail.hzau.edu.cn

¹ Biomass and Bioenergy Research Centre, Huazhong Agricultural University, Wuhan, China

Full list of author information is available at the end of the article

Background

Crop biomass residues represent enormous lignocellulose resource for the production of biofuels and chemicals [1]. Currently, lignocellulosic ethanol is considered as a promising short-term alternative to fossil fuels, due to its abundance in the cropping systems and having no conflict between energy demand and food supply [2, 3]. The production of lignocellulosic ethanol involves three major steps, including pretreatments for cell wall destruction, enzymatic hydrolysis for sugar release, and yeast fermentation for ethanol production [4, 5]. Because lignocellulose recalcitrance is a major hindrance for biomass conversion, genetic modification of plant cell walls has been proposed for enhancing biomass saccharification. However, plant cell walls have complicated structures and diverse functions, and their modification may thus affect plant growth and mechanical strength, in particular on plant lodging resistance. Since lodging is a major integrated trait tightly associated with grain and biomass yields in cereal crops [6–8], it becomes important to enhance both biomass saccharification and lodging resistance in transgenic plants.

Plant cell walls are mainly composed of cellulose, hemicelluloses, and lignin with small amounts of pectin and wall proteins. In principle, the wall polymers form a complex crosslink network with recalcitrant property against biomass saccharification [9–11]. In addition, the quantity and features of wall polymers are closely associated with plant lodging resistance [12–18]. For instance, cellulose crystallinity has been reported as the key factor that negatively affects either the biomass enzymatic digestibility or the plant lodging resistance [19]. In addition, the hemicelluloses' contents have been shown to negatively affect cellulose crystallinity, probably due to their association with cellulose microfibrils via hydrogen bonds [5, 19–21]. Hence, genetic modification of plant cell walls may be a promising solution to both biomass recalcitrance and plant lodging [9, 22–25].

Cellulose is the major component of plant cell walls and provides fermentable glucose for bioethanol production. In plants, cellulose synthase (CESA) catalyzes cellulose biosynthesis using UDP-glucose (UDPG) as substrate [26]. Sucrose synthase (SUS) catalyzes reversible conversion of sucrose and UDP into UDPG and fructose [27–29]. Importantly, SUS has been characterized to regulate cellulose biosynthesis by providing UDPG substrates in cotton fibers [30]. Although *SUS* transgenic plants have reportedly exhibited improvement in the cellulose-based traits [30–35], little is known about SUS impact on plant lodging resistance.

Rice is a major food crop around world with large amounts of biomass residues for biofuel production. In this study, we selected transgenic rice plants that

overexpressed *OsSUS3* gene using *Arabidopsis* cellulose synthase (*AtCesA8*) gene's promoter, specific for secondary cell wall synthesis. We detected biomass enzymatic saccharification and plant lodging resistance in the *AtCESA8::SUS3* transgenic rice plants. Furthermore, we detected any alterations of cell wall compositions and wall polymers features in the transgenic plants, and proposed a model that shows how *OsSUS3* overexpression leads to largely improved biomass enzymatic saccharification and plant lodging resistance by altering cellulose crystallinity and cell wall thickness.

Results


OsSUS3-transgenic plants maintain a normal growth with slightly increased biomass yields

Based on the public microarray database, we initially observed a low expression of *OsSUS3* gene in the tissues that are rich at secondary cell walls in rice (Additional file 1: Figure S1). To enhance *OsSUS3* expression, we selected the transgenic rice plants that expressed *OsSUS3* driven by *AtCesA8* promoter (Fig. 1a), which mainly drives cellulose synthase (*CESA*) gene's expression in the secondary cell wall synthesis of *Arabidopsis* [26]. As a result, the selected four independent homozygous lines were examined with much higher *OsSUS3* transcript levels in the stem tissues of transgenic rice plants, compared with the controls including 'Zhonghua11' (ZH11) cultivar and the transgenic line-expressed empty vector (EV) (Fig. 1b), which was confirmed by Western analysis (data not shown).

During the 2-year field experiments, all transgenic rice lines showed a normal growth and development over the life cycles of rice, compared with the ZH11 and EV line (Fig. 1c). However, the transgenic rice plants achieved slightly increased biomass yields by 6–8% at $P < 0.05$ and 0.01 levels (Fig. 1d; Additional file 1: Table S1), and the biomass yields were even consistent in two-year field experiments, based on a significantly positive correlation (Fig. 1e). The results suggest that overexpression of *OsSUS3* has little effect on plant growth with the slightly increased biomass yields in transgenic rice plants.

Enhanced biomass saccharification and bioethanol production in the *OsSUS3*-transgenic lines

In this study, biomass enzymatic saccharification (digestibility) was assessed by calculating the hexose yields released from enzymatic hydrolysis after chemical pretreatments of biomass residues in the mature stem tissues of rice (Fig. 2a). In general, all the four *AtCesA8::SUS3* transgenic lines exhibited significantly higher biomass saccharification than those of ZH11 and EV at $P < 0.01$ levels, with the hexoses yields being increased by 23–33 and 29–49% (per plant) or by 15–23 and 21–39% (per



dry matter) released from enzymatic hydrolysis after 1% NaOH and 1% H₂SO₄ pretreatments, respectively (Fig. 2b; Additional file 1: Figure S2a). Consequently, the *AtCESA8::SUS3* transgenic plants had the bioethanol yields increased by 24–47 and 44–63% (per plant) or by 13–37 and 29–54% (per dry matter), respectively, obtained from yeast fermentation (Fig. 2c; Additional file 1: Figure S2b). Hence, the overexpression of *OsSUS3* led to largely enhanced biomass saccharification and bioethanol production in rice.

Increased lodging resistance and mechanical strength in the *OsSUS3*-transgenic lines

Three criteria have been defined for evaluating plant lodging resistance. In particular, lodging index (LI) is highly related to plant height, fresh weight, and breaking force, which was the direct criterion that negatively accounts for plant lodging resistance [21, 36]. Extension and pushing forces are the other criteria accounting for the elasticity and mechanical strength of plant organs and tissues [8, 36–39]. In this study, we detected lodging index of the transgenic lines and controls from the two-year field experiments. All the four independent

transgenic lines exhibited reduced lodging index values by 17–50%, compared to those in ZH11 and EV. Meanwhile, we examined significantly increased extension force by 8–17% and pushing force by 13–44% in the *AtCESA8::SUS3* transgenic lines (Table 1). Because plant extension and pushing forces are tightly associated with lodging resistance [8, 36–39], all data were thus consistent with the findings about the significantly increased lodging resistance in the transgenic rice plants.

Altered cell wall ultrastructure and composition in the *OsSUS3*-transgenic lines

To understand enhanced biomass digestibility and lodging resistance of the *OsSUS3*-transgenic plants, we examined cell wall ultrastructure using scanning electron microscopy and transmission electron microscopy (Fig. 3). As a result, two representative *AtCESA8::SUS3* transgenic lines obviously exhibited thickened sclerenchyma cells (SC), vascular bundle cells (VB), and parenchyma cells (PC), compared to those in ZH11 and EV (Fig. 3a, b). In particular, the widths of entire cell walls and secondary cell walls in the sclerenchyma cells were respectively increased by 75–77 and 83–90% in the





Table 1 Detection of lodging index, extension, and pushing forces of the *OsSUS3*-transgenic lines in field experiments

Transgenic lines	Lodging index		Extension force (N)				Pushing force (N)			
	2013	2015	2013	2015	2015	2015				
Vector										
ZH11	170.62 ± 7.66	269.21 ± 8.47	173.80 ± 8.92	202.93 ± 7.18	1.66 ± 0.13					
EV	166.70 ± 6.34	278.64 ± 16.1	178.96 ± 4.28	206.78 ± 6.97	1.69 ± 0.14					
<i>AtCesA8::SUS3</i>										
1	108.14 ± 6.13**	-37% ^a	221.18 ± 7.59**	-18%	190.78 ± 3.08*	+10%	230.42 ± 4.79**	+14%	2.31 ± 0.10**	+39%
2	108.96 ± 7.82**	-36%	188.84 ± 22.72**	-30%	202.67 ± 15.59**	+17%	222.23 ± 3.54**	+10%	2.01 ± 0.06**	+21%
3	97.72 ± 8.57**	-43%	211.07 ± 11.23**	-22%	193.80 ± 11.04*	+12%	219.32 ± 3.67**	+8%	1.88 ± 0.09**	+13%
4	85.62 ± 9.24**	-50%	223.44 ± 14.06**	-17%	196.26 ± 11.28**	+13%	224.85 ± 12.46**	+11%	2.39 ± 0.10**	+44%

* and **, indicated significant difference between transgenic lines and ZH11 control by *t* test as *P* < 0.05 and 0.01 (*n* = 10)

^a Percentage of increased or decreased level between transgenic line and ZH11 by subtraction of two values divided by ZH11




AtCesA8::SUS3 transgenic plants, compared with the controls (Fig. 3c). Hence, the overexpression of *OsSUS3* led to remarkably increased secondary cell wall thickness in the transgenic rice plants.

Furthermore, we measured cell wall compositions of the *AtCesA8::SUS3* transgenic plants including cellulose, hemicelluloses, and lignin. As a comparison, the four transgenic lines had significantly increased cellulose levels at *P* < 0.05 and 0.01 levels (Fig. 3f; Additional file 1: Table S2), consistent with the Calcofluor staining specific for cellulose in the stem tissues (Fig. 3d). Upon using the plant glycan-directed monoclonal antibody (CCRC-M147), we observed that the *AtCesA8::SUS3* transgenic plants showed much brighter fluorescent signals specific for unsubstituted xylan of hemicelluloses in the thickened sclerenchyma cells and vascular bundle cells (Fig. 3e), consistent with the significantly increased

hemicelluloses levels by 9–21% in the transgenic lines (Fig. 3f; Additional file 1: Table S2). Meanwhile, we demonstrated that the four *AtCesA8::SUS3* transgenic lines had lignin levels similar to those of ZH11 and EV in the mature stems (Fig. 3f; Additional file 1: Table S2). Taking all these findings together, it is concluded that the overexpression of *OsSUS3* led to enhanced cellulose and hemicelluloses depositions into the cell walls, consistent with the findings of the more-thickened cell walls and increased biomass yields in transgenic rice plants.

Reduced cellulose crystallinity for enhanced biomass saccharification and lodging resistance

Cellulose crystallinity has been well characterized as the key negative factor on biomass enzymatic digestibility in different plant species [5, 19, 20]. Using X-ray detection method, this study measured cellulose crystalline index




(CrI) of the *AtCesA8::SUS3* transgenic plants, which is accounting for cellulose crystallinity [40–42]. As a comparison, all four transgenic rice lines had much lower CrI values by 7–10% than those of ZH11 and EV in the mature stem tissues (Fig. 4a). Because correlation analysis has been well applied to account for wall polymer feature impacts on biomass enzymatic saccharification in different plant species [5, 19, 20], we performed a correlation analysis between cellulose CrI and biomass saccharification or bioethanol production in all transgenic lines and controls. As a result, cellulose CrI values were negatively correlated with either the hexoses yields released from enzymatic hydrolysis after chemical pretreatments or the bioethanol yields from yeast fermentations at $P < 0.05$ or 0.01 levels, with high R^2 values at 0.498–0.657 (Fig. 4b, c). These data thus suggest that the reduction of cellulose CrI should be a major influential factor on the enhanced biomass saccharification and bioethanol production in the *AtCesA8::SUS3* transgenic plants, consistent with previous reports in other biomass samples [19, 40, 41]. Notably, the cellulose CrI also showed a significantly positive correlation with lodging index at $P < 0.01$ levels,

with high R^2 values at 0.705 (Fig. 4d), indicating that the reduced cellulose CrI should also be a major influential factor on the increased lodging resistance in transgenic rice plants. In addition, total hemicelluloses and cellulose levels exhibited a negative correlation with cellulose CrI values at $P < 0.01$ and 0.05 levels (Fig. 4e, f), consistent with the previous reports of hemicelluloses and cellulose having negative impacts on cellulose crystallinity [19].

Altered gene expressions involved in wall polymer synthesis

Because cellulose and hemicelluloses contents are increased in the *AtCesA8::SUS3* transgenic rice plants, we examined transcript levels of the genes that are involved in two wall polymers syntheses. As *OsCesA1*, 3, 8 and *OsCesA4*, 7, 9 are, respectively, involved in the cellulose synthesis of the primary and secondary cell walls in rice [42, 43], we demonstrated that those six *OsCesAs* genes had much higher expressions in the representative *AtCesA8::SUS3* transgenic plants at $P < 0.01$ level, compared to their expressions in the ZH11 control (Fig. 5a, b), which is consistent with the increased cellulose levels



in the transgenic plants. The *AtCesA8::SUS3* transgenic plant also showed much higher transcript levels of *OsIRX9* and *OsIRX14* than those of the ZH11 control at $P < 0.01$ levels (Fig. 5c). Because *OsIRX9* and *OsIRX14* have been identified to catalyze xylan backbone synthesis in rice [44, 45], the results were consistent with the increased hemicelluloses levels in the *AtCesA8::SUS3* transgenic plants.

Furthermore, we examined transcript levels of three representative *OsGH9B* genes in the *AtCesA8::SUS3* transgenic plants, because those three enzymes may have activity for reducing cellulose crystallinity in rice [19, 46]. Significantly, the *AtCesA8::SUS3* transgenic plant exhibited much higher *OsGH9B* expression levels than those of the ZH11 control (Fig. 5d), consistent with the findings of reduced cellulose CrI in the transgenic rice plants.


Discussion

Plant cell walls represent the most abundant renewable biomass resources for bioethanol production [1, 47] and largely affect mechanical strength of plants [21, 39, 48]. It has been characterized that plant cell wall compositions and wall polymer features play distinctive roles in biomass recalcitrance and plant lodging resistance [12–21]. Although genetic modification of plant cell walls has been posed as a promising solution to the recalcitrance and lodging [22–25], it remains a challenge to maintain normal growth and mechanical strength in transgenic plants. It thus becomes crucial to identify critical genes and appropriate promoters for selection of transgenic plants that are of improved biomass saccharification and lodging resistance [47]. In this study, the *AtCesA8::SUS3* transgenic rice plants could not only maintain a normal growth with slightly increased biomass yield, but also exhibited significantly enhanced biomass saccharification and lodging resistance, indicating a powerful genetic

strategy for cell wall modification in bioenergy crops. Although *SUS* transgenic plants have previously shown improvements in cellulose and starch contents in poplar, cotton, and potato [27–35], this study for the first time demonstrated the enhancement role of *SUS* in plant lodging resistance, which is one of the most important agronomic traits tightly associated with grain and biomass yields in crops.

To understand why the *AtCesA8::SUS3* transgenic rice plants are of improved saccharification and lodging resistance, we propose a mechanistic model to elucidate that the enhanced *OsSUS3* activity could provide sufficient UDPG substrate for direct cellulose biosynthesis and indirect hemicelluloses production, leading to reduced cellulose crystallinity and increased cell wall thickness in the transgenic rice plants (Fig. 6). The model hence highlights two major wall factors (cellulose crystallinity and cell wall thickening) that distinctively affect biomass enzymatic saccharification and plant lodging resistance in the transgenic rice plants. The cellulose crystallinity could negatively affect both biomass saccharification and plant lodging resistance, whereas the cell wall thickening mainly increase lodging resistance. Furthermore, it assumes that cellulose crystallinity should be negatively affected by both the cellulose modification with GH9B enzymes and the hemicelluloses deposition into the transgenic rice plants. The model also indicates that the syntheses of increased wall polymers (cellulose and hemicelluloses) with CESAs and IRXs enzymes should be the major factors of influence for the cell wall thickening in the transgenic plants.

It has been characterized that the cellulose CrI values reflect the relative amounts of crystalline materials and that the cellulose microfibrils with high crystallinity is thus less accessible by cellulases in biomass enzymatic hydrolysis [49]. On the other hand, the reduced cellulose



CrI may reflect an increased amorphous density of cellulose microfibrils that allows for more cellulase enzyme loading and facile accessibility, and this study has thus demonstrated that cellulose crystallinity is the key factor that negatively affects biomass enzymatic digestibility and bioethanol production in the *AtCESA8::SUS3* transgenic plants. In terms of the reduced cellulose crystallinity in the *AtCESA8::SUS3* transgenic plants, this study has provided two evidences: increased expressions of three representative *OsGH9B* genes and an increased hemicelluloses level, which has previously been reported to negatively affect cellulose crystallinity in rice and other plants [19, 46, 50, 51]. However, it remains to analyze cell wall and cellulose ultrastructures in the *AtCESA8::SUS3* transgenic plants in the future, which may provide a direct evidence about the reduced cellulose crystallinity.

As plant cell walls basically determine plants’ mechanical strength [12–19, 52], the increased cell wall thickness should be a major influential factor for the enhanced lodging resistance in the transgenic rice plants. Meanwhile, the increased extension and pushing forces should be also due to the cell wall thickenings in the transgenic plants, which is another evidence for the enhanced lodging resistance [36]. Because cellulose crystallinity negatively affects lodging index in rice, the reduction of cellulose crystallinity should be an additional contributor to the lodging resistance in the transgenic plants.

In addition, although the *OsSUS3*-transgenic plants had slightly increased biomass yield, they may have largely enhanced biomass production if grown in the large fields fed with good amounts of fertilizers in the future, because this study has witnessed a very high increase in cellulose and hemicelluloses levels and cell wall thickness. The previous report has indicated that the

SUS transgenic tree plants have a much enhanced biomass productivity [32, 33]. Therefore, the enhancement of cellulose accumulation in raw biomass and improvement of biomass digestibility will potentially reduce the costs of lignocellulosic biofuel productions [53]. In addition, those four independent *SUS3* transgenic lines displayed significantly decreased sucrose levels (Additional file 1: Table S3), suggesting that the improvements in the trait of transgenic plants are most likely due to an efficient energy allocation and consumption, particularly for cellulose synthesis.

Conclusions

Overexpression of *OsSUS3* driven by *AtCESA8* promoter could not only lead to enhanced biomass saccharification and bioethanol production after chemical pretreatments, but it also caused improved lodging resistance in the transgenic rice plants. Furthermore, this study has indicated that the reduction of cellulose crystallinity is a major factor for enhancements of both biomass saccharification and lodging resistance, and the increase of wall thickness may be the additional contributor to mechanical strength and lodging resistance in the transgenic rice plants. Hence, the results provide new insights into sucrose synthase regulation on carbon partitioning for cellulose and other polysaccharide biosynthesis. It also provides a powerful genetic approach for cell wall modification in bioenergy crops.

Methods

Vector constructs and transformation

The full-length cDNA of *SUS3* was cloned from rice cultivar ‘Nipponbare’ (a *japonica* variety) and sequenced. It was inserted into the plant binary vector pCAMBIA1300

(Cambia) under the *Arabidopsis* promoter AtCESA8. The constructs were introduced into *Agrobacterium tumefaciens* strain EHA105 and transferred to rice cultivar 'Zhonghua11' (ZH11) by *Agrobacterium*-mediated transformation. Ten sibling transgenic plants were assayed at each of the T₂–T₄ generations. All primers used for gene cloning are listed in Additional file 1: Table S4.

Total RNA isolation and RT-PCR analysis

Samples were collected from stems at heading stage. Total RNAs were extracted using Trizol reagent (Invitrogen, Carlsbad, CA, USA) and reverse-transcribed into cDNA with the GoScript™ Reverse Transcription System (Promega, USA). The RT-PCR reaction was performed as described previously [43]. Quantitative real time-PCR (qRT-PCR) was independently performed in triplicate using the SYBR Green PCR Master Mixture (ZF101, ZOMANBIO). A rice *polyubiquitin* gene (*OsUBQ1*) was used as the internal control. All primers used for RT-PCR are listed in Additional file 1: Table S4.

Field experiments and trait measurement

Transgenic rice plants were grown under natural field conditions in the experimental stations of Huazhong Agricultural University, Wuhan, China. Conventional rice cropping practices, including irrigation, fertilizer application, and pest control, were applied to the field experiments in this study.

Plant lodging index and mechanical property measurement

Plant lodging index (LI), extension force (EF) and pushing force (PF) were measured as described previously [19, 21, 36, 39]. All experiments were conducted on duplicate samples of ten sibling transgenic plants at each line, using the stem tissues at 30 days after flowering. The breaking resistance of the third internode was detected using a Prostrate Tester (DIK 7401, Daiki Rika Kogyo Co. Ltd., Tokyo, Japan), with the distance between fulcra of the tester at 5 cm. Fresh weight (*W*) of the upper portion of the plant was measured including panicle and the three internodes, leaf and leaf sheath. Bending moment (BM) and lodging index (LI) were calculated using the following formula: $BM = \text{Length from the third internode to the top of panicle} \times W$, $LI = BM / \text{breaking resistance}$. The EF were the stretching force of the samples before being broken, measuring with a universal force/length testing device (model RH-K300, Guangzhou, China). The PF were measured with five stems in each experimental unit (plot) by the prostrate tester (DIK-7400, Japan). The newton is used as the unit of EF and PF.

Plant cell wall fractionation and determination

The plant tissues were dried to constant weight, ground using a knife-mill and passed through a 40-mesh screen. Plant cell wall fractionation method and total cellulose and hemicelluloses assay were conducted as described previously [19, 21]. For crystalline cellulose extraction, biomass samples (0.1 g) were added with 5.0 mL acetic acid–nitric acid–water (8:1:2, v/v/v) and heated for 1 h in a boiling water bath with stirring every 10 min. After centrifugation, the pellet was washed several times with 5.0 mL water and dissolved in 67% H₂SO₄. Total hexoses in 67% H₂SO₄ were regarded as cellulose. For hemicelluloses analysis, the dry biomass powder samples (0.1 g) were treated by potassium phosphate buffer (pH 7.0), chloroform–methanol (1:1, v/v), DMSO–water (9:1, v/v), and 0.5% (w/v) ammonium oxalate to remove soluble sugar, lipids, starch, and pectin. The remaining pellet was suspended in 4 M KOH containing 1.0 mg mL⁻¹ sodium borohydride for 1 h at 25 °C, and the combined supernatant was neutralized, dialyzed and lyophilized for total hemicelluloses analysis.

Total lignin content includes acid-insoluble and -soluble lignin was determined by two-step acid hydrolysis method as previously described [54]. The sample (0.5 g, W1) was extracted with benzene–ethanol in a Soxhlet for 4 h, and then air dried. The sample was hydrolyzed with 10 mL 72% H₂SO₄ (v/v) in shaker at 30 °C for 1.5 h. After hydrolysis, the acid was diluted to a concentration of 2.88%, and then placed in the autoclave for 1 h at 121 °C. The autoclaved hydrolysis solution was vacuum-filtered through the filtering crucible. The filtrate was captured in a filtering flask for acid-soluble lignin. The acid-soluble lignin was solubilized during the hydrolysis process, and was measured by UV spectroscopy. The acid-insoluble residue was washed free of acid with distilled water and dried in an oven until attaining a constant weight. The weights of the crucible and dry residue were recorded (W2). Finally, the dried residues were ashed in the muffle furnace at 200 °C for 30 min and 575 °C for 4 h. The crucibles and ash were weighed, and their weights were recorded (W3). Acid-insoluble lignin (AIL) on original sample was calculated as follows: $AIL (\%) = (W2 - W3) \times 100 / W1\%$. Total lignin (%) = ASL% + AIL%. All experiments were conducted using biological triplicates.

Cellulose CrI detection

The X-ray diffraction (XRD) method was applied for detection of the lignocellulose crystalline index (CrI) in the crude cell wall materials using Rigaku-D/MAX instrument (Ultima III; Japan) as described by [46, 55–57]. The raw biomass powder was laid on the

glass sample holder ($35 \times 50 \times 5$ mm) and detected under plateau conditions. Ni-filtered Cu-K α radiation ($\lambda = 0.154056$ nm) was generated at voltage of 40 kV and current of 18 mA, and scanned at speed of $0.0197^\circ/\text{s}$ from 10 to 45° . The CrI was estimated using the intensity of the 200 peak (I_{200} , $\theta = 22.5^\circ$) and the intensity at the minimum between the 200 and 110 peaks (I_{am} , $\theta = 18.5^\circ$) as follows: $\text{CrI} = 100 \times (I_{200} - I_{\text{am}})/I_{200}$. The XRD method was detected with representative samples in triplicate.

Biomass pretreatment and enzymatic hydrolysis

The chemical (NaOH, H₂SO₄) pretreatment and sequential enzymatic hydrolysis were performed as described previously with minor modifications [57]. For NaOH pretreatment, the ground biomass powder was supplemented with 6 mL 1% NaOH (w/v). For H₂SO₄ pretreatment, the biomass powder was added with 6 mL 1% H₂SO₄ (v/v) and heated at 121 °C for 20 min in autoclave. The samples of chemical (NaOH, H₂SO₄) pretreatments were shaken at 150 r/min for 2 h at 50 °C, and centrifuged at 3000g for 5 min. The remaining pellet was washed three times with 10 mL distilled water, and all supernatants were combined for sugar analysis. The remaining residue was collected for enzymatic hydrolysis. Samples with 6 mL distilled water only were shaken for 2 h at 50 °C as control. All experiments were performed in the biological triplicates.

Yeast fermentation and ethanol measurement

Yeast fermentation and ethanol measurement were performed as previously described by Jin et al. [58] with minor modification. After pretreatments, the biomass residues and supernatants were neutralized to pH 4.8 using appropriate amounts of H₂SO₄ or NaOH. Then, mixed-cellulases were added to the final enzyme concentration at 1.6 g/L, and incubated at 150 r/min for 48 h at 50 °C. *Saccharomyces cerevisiae* (Angel yeast Co., Ltd., Yichang, China) was used in all fermentation reactions, and the yeast powder was dissolved in 0.2 M phosphate buffer (pH 4.8) for 30 min for activation prior to use. The activated yeast (ferment hexoses only) was inoculated into the mixture of enzymatic hydrolysates and residues with initial cell mass concentration at 0.5 g/L. The fermentation experiments were performed at 37 °C for 48 h, and distilled for determination of ethanol content. Ethanol content was measured using the dichromate oxidation method. All experiments were performed in the biological triplicates.

Scanning electron microscopy and transmission electron microscopy analyses

The second internodes (0.5 cm sections above the node) at the heading stage were cut into 1–2 mm pieces,

subsequently fixed with 2.5% (v/v) glutaraldehyde, vacuumed three times, and fixed for at least 24 h. Samples were natural dried, sputter-coated with gold particles, observed, and photographed using a scanning electron microscope (JSM-6390LV; JEOL, Japan). Scanning electron microscopy (SEM) analysis was based on at least three biological replications of the mounted specimens. All procedures were carried out according to the manufacturer's protocol.

Transmission electron microscopy (TEM) was used to observe cell wall structures in the middle 0.5 cm sections from the third leaf veins of three-leaves-old plants. Tissues were high-pressure frozen, freeze substituted, embedded, sectioned, and viewed according to McFarlane et al. [59]. The samples were post-fixed in 2% (w/v) OsO₄ for 1 h after extensively washing in the PBS buffer and embedded with Supr Kit (Sigma). Sample sections were cut with an Ultracut E ultramicrotome (Leica) and picked up on formvar-coated copper grids. After post-staining with uranyl acetate and lead citrate, the specimen were viewed under a Hitachi H7650 (Hitachi Ltd., Tokyo, Japan) transmission electron microscope. The width of cell wall was measured using the software ImageJ (NIH, USA), and more than 40 cell walls each for the different genotypes were measured. Significance was estimated using Student's *t* test.

Microscope observation

The sample preparation was performed as previously described [60]. The second internodes (0.5 cm sections above the node) at the heading stage were cut into pieces, subsequently fixed with 4% (w/v) paraformaldehyde, and dehydrated through an ethanol gradient (30, 50, 70, 90, 100 and 100%, each for 30 min), and then embedded in paraplast plus. The sections (8 μm thickness) were cut using a microtome (RM2265, Leica) and placed on lysine-treated slides which were dried for 2 days at 37 °C, and de-waxed with xylene and hydrated through an ethanol series (100–0%). The sections were treated with PBS buffer contained 3% SMP (skim milk powder, w/v) for 1 h, and incubated with PBS containing 10 $\mu\text{g}/\text{mL}$ CCRC-147 (<http://glycomics.ccruc.uga.edu/wall2/antibodies/antibodyHome.html>, CCRC-M147 recognized selectively all structures containing an unsubstituted xylan disaccharide) [61] for another 1 h. The immunolabeled samples were washed three times (5 min each) with PBS and incubated with a 100-fold dilution of anti-mouse-IgG in dark for 2 h. The anti-mouse-IgG antibody was labeled by fluorescein-isothiocyanate (FITC). Counterstaining was performed with Calcofluor white M2R fluorochrome (fluorescent brightener 28; Sigma; 0.25 $\mu\text{g}/\text{mL}$ in dH₂O). Immunofluorescence sections were imaged using a microscope (Olympus BX-61, Japan) equipped with

the following filter sets: 350/450 nm (ex/em) for visualizing calcofluor white stained cell walls, and 490/520 nm (ex/em) for green emission of the FITC fluorochrome, respectively.

Statistical calculation of correlation coefficients

Both two-tailed Student's *t* test and analysis of variance (ANOVA) were performed using SPSS. Significance was measured at the levels of $P < 0.05$ and $P < 0.01$. Correlation coefficients were calculated by performing Spearman's rank correlation analysis for all pairs of measured traits across the whole population.

Additional file

Additional file 1: Figure S1. Gene expression profiling of *OsSUS3* in life cycle of rice. **Figure S2.** Biomass enzymatic saccharification and ethanol production of the *OsSUS3*-transgenic rice plants. (a) Hexose yields released from enzymatic hydrolysis after the pretreatment with 1% NaOH or 1% H_2SO_4 . (b) Bioethanol yields obtained from yeast fermentation using the sugars released from biomass enzymatic hydrolysis as performed in (a). All data are given as means \pm SD. A Student's *t*-test was performed between transgenic plants and ZH11 as $^{**}P < 0.01$ and $^{*}P < 0.05$ ($n = 3$).

Abbreviations

SUS: sucrose synthase; UDPG: UDP-glucose; CrI: cellulose crystallinity index; Li: lodging index; EF: extension force; PF: pushing force; SEM: scanning electron microscopy; TEM: transmission electron microscopy; SC: sclerenchyma cells; VB: vascular bundle cells; PC: parenchyma cells; CESA: cellulose synthase; GH: glycoside hydrolase.

Authors' contributions

CF completed major experiments and prepared the manuscript; SF, JH, and YW participated in the cell wall determination experiments; LW, XL, and LW participated in lodging experiments; YT, TX, JL, and XC participated in enzymatic hydrolysis experiments; LP designed project, supervised experiments, and finalized the manuscript. All the authors read and approved the final manuscript.

Author details

¹ Biomass and Bioenergy Research Centre, Huazhong Agricultural University, Wuhan, China. ² National Key Laboratory of Crop Genetic Improvement, Huazhong Agricultural University, Wuhan, China. ³ College of Plant Science and Technology, Huazhong Agricultural University, Wuhan, China. ⁴ College of Life Science and Technology, Huazhong Agricultural University, Wuhan, China. ⁵ HaiKou Experimental Station, Chinese Academy of Tropical Agricultural Sciences, Haikou 570102, China. ⁶ Department of Plant Science, North Dakota State University, Fargo, ND, USA.

Acknowledgements

This work was supported in part by grants from the Fundamental Research Funds for the Central Universities of China (Program Nos. 2662015PY018; 2662015PY173), the National Transgenic Project (2009ZX08009-119B), the National 111 Project (B08032), the National Science Foundation of China (31670296; 31571721), and the Huazhong Agricultural University Scientific & Technological Self-innovation Foundation (2011SC01).

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 28 June 2017 Accepted: 8 September 2017

Published online: 16 September 2017

References

- Service RF. Biofuel researchers prepare to reap a new harvest. *Science*. 2007;315(5818):1488–91.
- Himmel ME, Bayer EA. Lignocellulose conversion to biofuels: current challenges, global perspectives. *Curr Opin Biotechnol*. 2009;20(3):316–7.
- Yuan JS, Tiller KH, Al-Ahmad H, Stewart NR, Stewart CN Jr. Plants to power: bioenergy to fuel the future. *Trends Plant Sci*. 2008;13(8):421–9.
- Carroll A, Somerville C. Cellulosic biofuels. *Annu Rev Plant Biol*. 2009;60:165–82.
- Xu N, Zhang W, Ren SF, Liu F, Zhao CQ, Liao HF, Xu ZD, Huang JF, Li Q, Tu YY, et al. Hemicelluloses negatively affect lignocellulose crystallinity for high biomass digestibility under NaOH and H_2SO_4 pretreatments in *Miscanthus*. *Biotechnol Biofuels*. 2012;5:58.
- Kono M. "Physiological aspects of lodging," in *Science of the Rice Plant*. In: Matsuo T, Kumazawa K, Ishii R, Ishihara K, Hirata H, editors. Physiology, vol. 2. Tokyo: Food and Agriculture Policy Research Center; 1995. p. 971–82.
- Weber CR, Fehr WR. Seed yield losses from lodging and combine harvesting in soybeans. *Agron J*. 1966;58(3):287–9.
- Berry PM, Sterling M, Spink JH, Baker CJ, Sylvester-Bradley R, Mooney SJ, Tams AR, Ennos AR. Understanding and reducing lodging in cereals. *Adv Agron*. 2004;84:217–71.
- Pauly M, Keegstra K. Plant cell wall polymers as precursors for biofuels. *Curr Opin Plant Biol*. 2010;13(3):305–12.
- Chen F, Dixon RA. Lignin modification improves fermentable sugar yields for biofuel production. *Nat Biotechnol*. 2007;25(7):759–61.
- Loque D, Scheller HV, Pauly M. Engineering of plant cell walls for enhanced biofuel production. *Curr Opin Plant Biol*. 2015;25:151–61.
- Islam MS, Peng S, Visperas RM, Ereful N, Bhuiya MSU, Julfikar AW. Lodging-related morphological traits of hybrid rice in a tropical irrigated ecosystem. *Field Crops Res*. 2007;101(2):240–8.
- Kelbert AJ, Spaner D, Briggs KG, King JR. The association of culm anatomy with lodging susceptibility in modern spring wheat genotypes. *Euphytica*. 2004;136(2):211–21.
- Tripathi SC, Sayre KD, Kaul JN, Narang RS. Growth and morphology of spring wheat (*Triticum aestivum* L.) culms and their association with lodging: effects of genotypes N levels and ethephon. *Field Crop Res*. 2003;84(3):271–90.
- Zuber U, Winzeler H, Messmer MM, Keller M, Keller B, Schmid JE, Stamo P. Morphological traits associated with lodging resistance of spring wheat (*Triticum aestivum* L.). *J Agron Crop Sci*. 1999;182(1):17–24.
- Ishimaru K, Togawa E, Ookawa T, Kashiwagi T, Madoka Y, Hirotsu N. New target for rice lodging resistance and its effect in a typhoon. *Planta*. 2008;227(3):601–9.
- Kashiwagi T, Ishimaru K. Identification and functional analysis of a locus for improvement of lodging resistance in rice. *Plant Physiol*. 2004;134(2):676–83.
- Kashiwagi T, Madoka Y, Hirotsu N, Ishimaru K. Locus *prl5* improves lodging resistance of rice by delaying senescence and increasing carbohydrate reaccumulation. *Plant Physiol Biochem*. 2006;44(2):152–7.
- Li FC, Zhang ML, Guo K, Hu Z, Zhang R, Feng YQ, Yi XY, Zou WH, Wang LQ, Wu CY, et al. High level hemicellulosic arabinose predominately affects lignocellulose crystallinity for genetically enhancing both plant lodging resistance and biomass enzymatic digestibility in rice mutants. *Plant Biotechnol J*. 2015;13(4):514–25.
- Li FC, Ren SF, Zhang W, Xu ZD, Xie GS, Chen Y, Tu YY, Li Q, Zhou SG, Li Y, et al. Arabinose substitution degree in xylan positively affects lignocellulose enzymatic digestibility after various NaOH/ H_2SO_4 pretreatments in *Miscanthus*. *Bioresour Technol*. 2013;130:629–37.
- Li FC, Xie GS, Huang JF, Zhang R, Li Y, Zhang MM, Wang YT, Li A, Li XK, Xia T, et al. OsCESA9 conserved-site mutation leads to largely enhanced plant lodging resistance and biomass enzymatic saccharification by reducing cellulose DP and crystallinity in rice. *Plant Biotech J*. 2017. doi:10.1111/pbi.12700.
- Demura T, Ye Z. Regulation of plant biomass production. *Curr Opin Plant Biol*. 2010;13(3):299–304.

23. Torney F, Moeller L, Scarpa A, Wang K. Genetic engineering approaches to improve bioethanol production from maize. *Curr Opin Plant Biol*. 2007;18(3):193–9.
24. Vega-Sánchez ME, Ronald PC. Genetic and biotechnological approaches for biofuel crop improvement. *Curr Opin Biotechnol*. 2010;21(2):218–24.
25. Xie GS, Peng LC. Genetic engineering of energy crops: a strategy for biofuel production in China. *J Integr Plant Biol*. 2011;53(2):143–50.
26. Taylor NG, Laurie S, Turner SR. Multiple cellulose synthase catalytic subunits are required for cellulose synthesis in *Arabidopsis*. *Plant Cell*. 2000;12(12):2529–40.
27. Sun J, Loboda T, Sung SJS, Black CC. Sucrose synthase in wild tomato, *Lycopersicon chmielewskii*, and tomato fruit sink strength. *Plant Physiol*. 1992;98(3):1163–9.
28. Amor Y, Haigler C, Johnson S, Wainscott M, Delmer DP. A membrane-associated form of sucrose synthase and its potential role in synthesis of cellulose and callose in plants. *Proc Natl Acad Sci USA*. 1995;92(20):9353–7.
29. Zrenner R, Salanoubat M, Willmitzer L, Sonnwald U. Evidence of the crucial role of sucrose synthase for sink strength using transgenic potato plants (*Solanum tuberosum* L.). *Plant J*. 1995;7(1):97–107.
30. Jiang Y, Guo W, Zhu H, Ruan YL, Zhang T. Overexpression of *GhSusA1* increases plant biomass and improves cotton fiber yield and quality. *Plant Biotechnol J*. 2012;10(3):301–12.
31. Konishi T, Ohmiya Y, Hayashi T. Evidence that sucrose loaded into the phloem of a poplar leaf is used directly by sucrose synthase associated with various β -glucan synthases in the stem. *Plant Physiol*. 2004;134(3):1146–52.
32. Coleman HD, Ellis DD, Gilbert M, Mansfield SD. Up-regulation of sucrose synthase and UDP-glucose pyrophosphorylase impacts plant growth and metabolism. *Plant Biotechnol J*. 2006;4(1):87–101.
33. Coleman HD, Yan J, Mansfield SD. Sucrose synthase affects carbon partitioning to increase cellulose production and altered cell wall ultrastructure. *Proc Natl Acad Sci USA*. 2009;106(31):13118–23.
34. Xu SM, Brill E, Llewellyn DJ, Furbank RT, Ruan YL. Overexpression of a potato sucrose synthase gene in cotton accelerates leaf expansion, reduces seed abortion, and enhances fiber production. *Mol Plant*. 2012;5(2):430–41.
35. Poovaiah CR, Mazarei M, Decker SR, Turner GB, Sykes RW, Davis MF, Stewart CN. Transgenic switchgrass (*Panicum virgatum* L.) biomass is increased by overexpression of switchgrass sucrose synthase (*PvSUS1*). *Biotechnol J*. 2015;10(4):552–63.
36. Fan CF, Li Y, Hu Z, Hu HZ, Wang GY, Li A, Wang YM, Tu YY, Xia T, Peng LC, et al. Ectopic expression of a novel OsExtensin-like gene consistently enhances plant lodging resistance by regulating cell elongation and cell wall thickening in rice. *Plant Biotech J*. 2017. doi:10.1111/pbi.12766.
37. Crook MJ, Ennos AR. Stem and root characteristics associated with lodging resistance in four winter wheat cultivars. *J Agric Sci*. 1994;123(2):167–74.
38. Hai L, Guo HJ, Xiao SH, Jiang GL, Zhang XY, Yan CS, Xin ZY, Jia JZ. Quantitative trait loci (QTL) of stem strength and related traits in a doubled-haploid population of wheat (*Triticum aestivum* L.). *Euphytica*. 2005;141(1):1–9.
39. Zhang ML, Wei F, Guo K, Hu Z, Li YY, Xie GS, Wang YT, Cai XW, Peng LC, Wang LQ. A novel *FC116/BC10* mutation distinctively causes alteration in the expression of the genes for cell wall polymer synthesis in rice. *Front Plant Sci*. 2016;7:1366.
40. Wu ZL, Zhang ML, Wang LQ, Tu YY, Zhang J, Xie GS, Zou WH, Li FC, Guo K, et al. Biomass digestibility is predominantly affected by three factors of wall polymer features distinctive in wheat accessions and rice mutants. *Biotechnol Biofuels*. 2013;6:183.
41. Zhang W, Yi ZL, Huang JF, Li FC, Hao B, Li M, Hong SF, Lv YZ, Sun W, et al. Three lignocellulose features that distinctively affect biomass enzymatic digestibility under NaOH and H₂SO₄ pretreatments in *Miscanthus*. *Bioresour Technol*. 2013;130:30–7.
42. Tanaka K, Murata K, Yamazaki M, Onosato K, Miyao A, Hirochika H. Three distinct rice cellulose synthase catalytic subunit genes required for cellulose synthesis in the secondary wall. *Plant Physiol*. 2003;133(1):73–83.
43. Wang LQ, Guo K, Li Y, Tu YY, Hu HZ, Wang BR, Cui XC, Peng LC. Expression profiling and integrative analysis of the *CESA/CSL* superfamily in rice. *BMC Plant Biol*. 2010;10(1):282.
44. Scheller HV, Ulvskov P. Hemicelluloses. *Annu Rev Plant Biol*. 2010;61:263–89.
45. Chiniquy D, Varanasi P, Oh T, Harholt J, Katnelson J, Singh S, Auer M, Simmons B, Adams PD, Scheller HV, et al. Three novel rice genes closely related to the *Arabidopsis irx9*, *irx9l*, and *irx14* genes and their roles in xylan biosynthesis. *Front Plant Sci*. 2013;4:83.
46. Xie GS, Yang B, Xu ZD, Li FC, Guo K, Zhang ML, Wang LQ, Zou WH, Wang YT, Peng LC. Global identification of multiple OsGH9 family members and their involvement in cellulose crystallinity modification in rice. *PLoS ONE*. 2013;8:e50171.
47. Wang YT, Fan CC, Hu HZ, Li Y, Sun D, Wang YM, Peng LC. Genetic modification of plant cell walls to enhance biomass yield and biofuel production in bioenergy crops. *Biotechnol Adv*. 2016;34(5):997–1017.
48. Hu KM, Cao JB, Zhang J, Xia F, Ke YG, Zhang HT, Xie WY, Liu HB, Cui Y, Cao YL, et al. Improvement of multiple agronomic traits by a disease resistance gene via cell wall reinforcement. *Nat Plants*. 2017;3:17009.
49. Himmel ME, Ding SY, Johnson DK, Adney WS, Nimlos MR, Brady JW, Foust TD. Biomass recalcitrance: engineering plants and enzymes for biofuels production. *Science*. 2007;315(5813):804–7.
50. Takahashi J, Rudsander UJ, Hedenström M, Banasiak A, Harholt J, Amelot N, Immerzeel P, Ryden P, Endo S, Ibatullin FM, et al. KORRIGAN1 and its aspen homolog PttCel9A1 decrease cellulose crystallinity in *Arabidopsis* stems. *Plant Cell Physiol*. 2009;50(6):1099–115.
51. Kalluri UC, Payyavula RS, Labbé JL, Engle N, Bali G, Jawdy SS, Sykes R, Davis M, Ragauskas A, Tuskan G, et al. Down-regulation of KORRIGAN-like endo- β -1,4-glucanase genes impacts carbon partitioning, mycorrhizal colonization and biomass production in *Populus*. *Front Plant Sci*. 2016;7:1455.
52. Somerville C. Cellulose synthesis in higher plants. *Annu Rev Plant Biol*. 2006;22:53–78.
53. Klein-Marcuschamer D, Oleskowicz-Popiel P, Simmons BA, Blanch HW. Technoeconomic analysis of biofuels: a wiki-based platform for lignocellulosic biorefineries. *Biomass Bioenergy*. 2010;34(12):1914–21.
54. Li M, Si S, Hao B, Zha Y, Wan C, Hong S, Kang Y, Jia J, Zhang J, Li M, et al. Mild alkali-pretreatment effectively extracts guaiacyl-rich lignin for high lignocellulose digestibility coupled with largely diminishing yeast fermentation inhibitors in *Miscanthus*. *Bioresour Technol*. 2014;169:447–54.
55. Sun D, Alam A, Tu YY, Zhou SG, Wang YT, Xia T, Huang JF, Ying L, Zahoor Wei XY, et al. Steam-exploded biomass saccharification is predominately affected by lignocellulose porosity and largely enhanced by Tween-80 in *Miscanthus*. *Bioresour Technol*. 2017;239:74–81.
56. Tu YY, Wang LQ, Xia T, Sun D, Zhou SG, Wang YT, Ying L, Zhang HP, Zhang T, et al. Mild chemical pretreatments are sufficient for complete saccharification of steam-exploded residues and high ethanol production in desirable wheat accessions. *Bioresour Technol*. 2017;243:319–26.
57. Sun D, Li Y, Wang J, Tu YY, Wang YT, Hu Z, Zhou SG, Wang LQ, Xie GS, et al. Biomass saccharification is largely enhanced by altering wall polymer features and reducing silicon accumulation in rice cultivars harvested from nitrogen fertilizer supply. *Bioresour Technol*. 2017;243:957–65.
58. Jin WX, Chen L, Hu M, Sun D, Ao L, Ying L, Zhen H, Zhou SG, Tu YY, Xia T, et al. Tween-80 is effective for enhancing steam-exploded biomass enzymatic saccharification and ethanol production by specifically lessening cellulase adsorption with lignin in common reed. *Appl Energy*. 2016;175:82–90.
59. McFarlane HE, Young RE, Wasteneys GO, Samuels AL. Cortical microtubules mark the mucilage secretion domain of the plasma membrane in *Arabidopsis* seed coat cells. *Planta*. 2008;227:1363–75.
60. Cao YP, Li JL, Yu L, Chai GH, He G, Hu RB, Qi G, Kong YZ, Fu CX, Zhou GK. Cell wall polysaccharide distribution in *Miscanthus lutarioriparius* stem using immuno-detection. *Plant Cell Rep*. 2014;33:643–53.
61. Schmidt D, Schuhmacher F, Geissner A, Seeberger PH, Pfrenge F. Automated synthesis of arabinoxylan-oligosaccharides enables characterization of antibodies that recognize plant cell wall glycans. *Chem Eur J*. 2015;21:5709–13.